

# **Flux and Wave Measurements during the Adverse Weather Experiment**

William M. Drennan  
Applied Marine Physics Division  
Rosenstiel School of Marine and Atmospheric Science  
University of Miami  
4600 Rickenbacker Causeway, Miami, FL 33149-1098  
phone:(305)-361-4798 fax:(305)-361-4701 email: [wdrennan@rsmas.miami.edu](mailto:wdrennan@rsmas.miami.edu)

Award Number: N000140010594  
<http://www.oe.fau.edu/~manhar/awe2000/>

## **LONG-TERM GOALS**

The long-term goal is to investigate the impact of an adverse weather front on mixing, air-sea interaction, and sediment transport in the shallow water column over the continental shelf.

## **OBJECTIVES**

The objectives of the Adverse Weather Experiment (AWE) are: i) to parameterize the vertical fluxes of heat and momentum and the turbulent kinetic energy (TKE) dissipation rate in the upper oceanic mixed layer, and ii) to parameterize the buoyancy and momentum fluxes in the marine atmospheric boundary layer (MABL) during the passage of an adverse weather front over the warm continental shelf, off the east coast of Florida. A related objective is to employ the new parameterizations to improve present, laboratory based, sub-grid closure schemes used in turbulence models of the oceanic boundary layer. This project is part of a multi-institution effort. The objectives of this component of the project are:

- A) To measure fluxes (momentum and heat) in the MABL, along with supporting wave (directional spectra) and meteorological parameters at a site adjacent to the AWE-AUV survey area;
- B) To use the above measurements, along with those of OSCAR and an ADCP to parameterize measured TKE dissipation rates by wind stress, current shear, buoyancy and wave parameters.

## **APPROACH**

The key variables thought to influence TKE dissipation rates in the upper ocean mixed layer are the wind stress, buoyancy flux (Anis and Moum, 1995), horizontal shear, and surface wave field (Terray et al, 1996). Hence measurement of these parameters is a very important part of an effort to parameterize TKE dissipation rates. During the Adverse Weather Experiment measurements were carried out as follows:

- Mobile autonomous underwater vehicles (AUVs) were utilized to make measurements of current profiles, conductivity, temperature and small scale turbulence, including TKE dissipation rates (Dhanak and Holappa, FAU).

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2001</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2001 to 00-00-2001</b>	
4. TITLE AND SUBTITLE <b>Flux and Wave Measurements during the Adverse Weather Experiment</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Applied Marine Physics Division,,Rosenstiel School of Marine and Atmospheric Science,,University of Miami,4600 Rickenbacker Causeway,,Miami,,FL, 33149</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <b>The long-term goal is to investigate the impact of an adverse weather front on mixing, air-sea interaction, and sediment transport in the shallow water column over the continental shelf.</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>6</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

- An air-sea interaction spar (ASIS) buoy was deployed adjacent to the AUV survey area to measure eddy-correlation momentum and buoyancy fluxes in the MABL, along with surface wave spectra and supporting mean meteorological and sea surface parameters (Drennan, RSMAS).
- An ocean surface current radar (OSCR) and a moored Acoustic Doppler Current Profiler (ADCP) were deployed continuously during AWE to measure the horizontal and vertical surface current structures, respectively, over the period of the experiment (Shay, RSMAS).


Using the data collected over the continental shelf during several adverse weather events, the idea is to parameterize the TKE dissipation rate in terms of the key forcing parameters.

## WORK COMPLETED

On 7 April 2000, an ASIS buoy was moored at 26° 3.75' N 80° 5.42' W, at 20 m depth, off Dania, FL, at the northern end of the AUV survey area. The buoy remained on site for 5 weeks, and was recovered on 11 May 2000. During the deployment, the buoy was instrumented for flux and wave measurements as follows: 3D wind vector (Solent 1012R3A sonic anemometer), air temperature (2 fast response thermistors, and Rotronic), humidity (Rotronic), surface elevation (8-gauge capacitance wave wire array), surface currents+SST+salinity (SensorTek UCM-60DL) and buoy motion (3 accelerometers, 3 rate gyros, 1 compass). During AWE the data return rate for most sensors was near 100% throughout the deployment period. The data have been processed to yield shear and buoyancy forcing, along with wave, surface current and mean meteorological information (Figure 1). Work is ongoing, in collaboration with other AWE participants, to address how these forcing parameters relate to larger scale features and oceanic turbulence.


## RESULTS

A summary of the ASIS data is shown in Figures 1 and 2. In Fig. 1, the wind speed, air temperature and sea surface temperature are shown, along with momentum and buoyancy forcing parameters (friction velocity,  $u_*$ , and Obukhov length,  $L$ ) derived from the measured turbulent fluxes. The Obukhov length, related to the ratio of buoyant to shear forcing, has been calculated following two approaches: one using a bulk buoyancy flux derived from measured mean temperatures and humidity, and the other using a buoyancy flux calculated directly from the sonic temperature flux (obtained via eddy correlation), with corrections following Dupuis et al. (1997). Both are plotted in Figure 1, and the agreement is very good, although there is evidently a systematic underestimate of bulk  $L$  compared with sonic-derived  $L$  in the unstable regime. This underestimate of roughly 20% is consistent with a reduction of the measured bulk SST by 0.7°C. During AWE, the bulk SST was measured at a depth of roughly 5m, and hence may not be representative of the near-surface layer during much of the day (e.g. Price et al. 1986; Fairall et al 1996).



**Figure 1: Conditions during AWE experiment. The panels show:**  
**(top) Obukhov length  $|L|$ :  $L<0$  -Blue;  $L>0$  -Red;  $L_{sonic}$  (solid);  $L_{bulk}$  (•);**  
**(middle) Air temperature -Blue, and sea surface temperature -Red;**  
**(bottom) Wind speed at 7m - Blue, and friction velocity - Red**  
*[Primary features of figure are described in main text]*

Of special interest for the scientific objectives are the two periods JD 99-100 and 109-110 when cold fronts passed through the domain – note the wind shifts and changes in air temperature in Fig. 1. Note also the rapid increase in wave energy during the first period as the wind shifts onshore (Fig. 2). These periods coincide with AUV missions (Dhanak, FAU) during which TKE dissipation rates were measured, and hence are of primary interest in achieving the experimental goals. OSCAR surface current data (Shay, RSMAS) are available throughout the ASIS deployment period. Another event of interest is JD 114, when cool continental air was advected over the warmer coastal waters, resulting in high buoyancy forcing.



**Figure 2: Wave conditions during AWE experiment. The panels show: top) Significant wave height; middle) Inverse wave age; bottom) Evolution of 1D surface elevation spectrum. The contours showing spectral energy are spaced logarithmically. [Primary features of figure are described in main text]**

## IMPACT/APPLICATIONS

These are the first simultaneous measurements of TKE dissipation rates along with all the principal forcing terms (wind shear, buoyancy flux, surface waves and horizontal shear). These measurements, taken in a variety of meteorological and wave conditions, with varying surface current shear, will allow us to parameterize the TKE dissipation rate in terms of the important physical forcing variables. This in turn will allow for improvements to the sub-grid closure schemes used in turbulence models of the upper ocean mixed layer.

## TRANSITIONS

The ASIS data are presently being used by other participants in the AWE experiment. Transitions to the wider community are expected upon completion of the joint analysis. Several manuscripts are in preparation.

## RELATED PROJECTS

The Adverse Weather Experiment is a multi-institution effort involving three ONR-funded projects. The other two projects are Dhanak and Holappa (FAU) and Shay (RSMAS). Lueck (U Victoria) is also a participant in AWE.

## REFERENCES

- Anis, A., and J.N. Moum, 1995: Surface wave-turbulence interactions: scaling  $\varepsilon(z)$  near the sea surface. *J. Phys. Oceanogr.* **25**, 2025-2045.
- Dupuis, H., P.K. Taylor, A. Weill and K. Katsaros, 1997: Inertial dissipation method applied to derive turbulent fluxes over the ocean during the Surface of the Ocean, Fluxes and Interactions with the Atmosphere/ Atlantic Stratocumulus Transition Experiment (SOFIA/ASTEX) and Structure des Echanges Mer-Atmosphere, Proprietes des Heterogeneites Oceaniques: Recherche Experimentale (SEMAPHORE) experiments with low to moderate wind speeds. *J. Geophys. Res.* **102**, 21115-21129.
- Fairall, C.W., E.F. Bradley, J.S. Godfrey, G.A. Wick, J.B. Edson and G.S. Young, 1996: Cool-skin and warm-layer effects on sea surface temperature. *J. Geophys. Res.*, **101**, 1295-1308.
- Price, J.F., R.A. Weller, and R. Pinkel, 1986: Diurnal cycling: Observations and models of the upper ocean response to diurnal heating, cooling and wind mixing. *J. Geophys. Res.*, **91**, 8411-8427.
- Terray, E.A., M.A. Donelan, Y.C. Agrawal, W.M. Drennan, K.K. Kahma, A.J. Williams III, P.A. Hwang and S.A. Kitaigorodskii, 1996: Estimates of kinetic energy dissipation under breaking waves. *J. Phys. Oceanogr.* **26**, 792-807.